ON GRAPHS UNIQUELY DEFINED BY THEIR K-CIRCULAR MATROIDS

José F. De Jesús

University of Puerto Rico, San Juan, Puerto Rico, United States

Alexander Kelmans

University of Puerto Rico, San Juan, Puerto Rico, United States
Rutgers University, New Brunswick, New Jersey, United States

arXiv:1508.07627v1 [math.CO] 30 Aug 2015

Abstract

In 30's Hassler Whitney considered and completely solved the problem (WP) of describing the classes of graphs G having the same cycle matroid $M(G)$ [14, 15]. A natural analog $(WP)'$ of Whitney's problem (WP) is to describe the classes of graphs G having the same matroid $M'(G)$, where $M'(G)$ is a matroid (on the edge set of G) distinct from $M(G)$. For example, the corresponding problem $(WP)' = (WP)_\theta$ for the so-called bicircular matroid $M_\theta(G)$ of graph G was solved in [2, 11]. In [6] we introduced and studied the so-called k-circular matroids $M_k(G)$ for every non-negative integer k that is a natural generalization of the cycle matroid $M(G) := M_0(G)$ and of the bicircular matroid $M_\theta(G) := M_1(G)$ of graph G. In this paper (which is a continuation of our paper [6]) we establish some properties of graphs guaranteeing that the graphs are uniquely defined by their k-circular matroids.

Key words: graph, vertex star, bicycle, cacti-graph, matroid, k-circular matroid, non-separating cocircuit.

MSC Subject Classification: 05B35, 05C99

1 Introduction

In 30's Hassler Whitney developed a remarkable theory on the matroid isomorphism and the matroid duality of graphs [13–16]. He considered a graph G and the so called cycle matroid $M(G)$ of G (whose circuits are the edge subsets of the cycles in G) and stated the following natural problems on pairs $\langle G, M(G)\rangle$:

(WP) describe the classes of graphs having the same cycle matroid and, in particular, graphs that can be reconstructed from cycle matroid (up to the names of vertices) and

(WP^*) describe the pairs of graphs whose cycle matroids are dual, i.e. describe the class of graphs closed under their cycle matroids duality.

Classical Whitney's graph matroid-isomorphism theorem and Whitney's planarity criterion provide the answers to the above questions [14–16] (see also [10]).

Naturally, Whitney's problems and interesting results along this line prompted further questions and research on possible strengthenings as well as various extensions or analogs of some Whitney's results (see, for example, [2, 4, 5, 7, 8, 11]).

In [6] we introduced and studied the so-called k-circular matroids $M_k(G)$ of graph G. It is natural to consider an analog $(WP)_k$ of Whitney's problem $(WP) := (WP)_0$ on the classes of graphs having the same k-circular matroid. In this paper (which is a continuation of our paper [6]) we consider a particular problem of $(WP)_k$ on graphs uniquely defined by their k-circular matroids.

Section 2 provides some notions, notation, and some necessary preliminary facts on matroids and graphs. In particular, we introduce a classification of the vertex stars of a graph G in terms of the corank of matroid $M_k(G)$. This classification will play a key role in the study of the problem on describing the graphs uniquely defined by their k-circular matroids.

In Section 3 we describe the vertex stars in a graph G that are cocircuits of matroid $M_k(G)$.

In Section 4 we characterize the so-called non-separating cocircuits of matroid $M_k(G)$ in terms of graph G. This notion will be essential in proving some results on pairs $\langle G, M_k(G)\rangle$ analogous to Whitney's matroid isomorphism theorem.

By Whitney's matroid isomorphism theorem a graph with at least 4 vertices is uniquely defined by its cycle matroid if and only if the graph is multi 3-connected. In Section 5 we provide an extension of above Whitney's result by describing a class of graphs uniquely defined by their k-circular matroid. Every vertex star in a graph G of this class is a non-separating cocircuit of $M_k(G)$. This result as well as the main result of the next section is based on the fact that if S is a non-separating cocircuit of $M_k(G)$, then S is not only a vertex star of G but also a vertex star of every graph G' with $M_k(G') = M_k(G)$.

In Section 6 we describe another class of graphs uniquely defined by their k-circular matroid. For each graph G in this class every vertex star of G except for one is a non-separating cocircuit of $M_k(G)$.

2 Main notions and notation

The basic notions and notation we use here is the same as in our paper [6] since this paper is a continuation of [6] (see also [1, 3, 10, 12]). We will remind here some of these notions and notation.

A graph G is called *cacti-graph* if G has no isolated vertices, no leaves, and no cycle components. A connected cacti-graph is called a *cactus*. Let $\mathcal{G}_{\bowtie}$ denote the set of cacti-graphs and $\mathcal{CG}_{\bowtie}$ denote the set of connected graphs from $\mathcal{G}_{\bowtie}$, and so each member of $\mathcal{CG}_{\bowtie}$ is a cactus.

Let $\mathcal{G}$ be the set of finite graphs and $G \in \mathcal{G}$. Let $\Delta(G) = |E(G)| - |V(G)|$.
Instead of $\Delta(G)$ we will write simply ΔG. Let $X \subseteq E(G)$ and $G\langle X\rangle$ be the subgraph of G induced by X. Then $\Delta(G\langle X\rangle) = |X| - |V(G\langle X\rangle)|$.

For a graph $G = (V, E, \phi)$ and $k \geq 1$, let $\mathcal{C}_k(G) = \{C \subseteq E : \Delta G\langle C\rangle = k \text{ and } G\langle C\rangle \in \mathcal{G}_{\bowtie}\}$. Then $\mathcal{C}_k(G)$ is the collection of circuits of a matroid on E (see [6]). We call $M_k(G) = (E, \mathcal{C}_k(G))$ the *k-circular matroid of graph G*.

Let $\mathcal{I}_k(G)$, $\mathcal{B}_k(G)$, and $\mathcal{C}_k^*(G)$ denote the families of independent sets, bases, and cocircuits of $M_k(G)$, respectively. Let $\rho_k(G)$ and $\rho_k^*(G)$ denote the rank and the corank of matroid $M_k(G)$.

For a graph G with at least one cycle, let $\lfloor G \rfloor$ denote the maximum subgraph of G with no leaves and no isolated vertices. Graph $\lfloor G \rfloor$ is called the *kernel of graph G*. If F is a forest, then the kernel of F is not defined.

For a graph G having a component with at least two cycles, let $[G]$ denote the maximum subgraph of G with no leaves, no isolated vertices, and no cycle components. Graph $[G]$ is called *the core of graph G*. If every component of G has at most one cycle, then the core of G is not defined.

We defined a matroid $M = (E, \mathcal{I})$ to be *connected* if M has no loops, no coloops and if every two elements in E belong to a common circuit of M.

We call a cocircuit C^* of a connected matroid M a *non-separating cocircuit* of M if $M \setminus C^*$ is a connected matroid (see [9]).

Here is some notation we will use:

$S(v, G)$ is the *v-star in G*, i.e. the set of edges incident to vertex v in G and $s(v, G) = |S(v, G)|$,

$\mathcal{S}(G) = \{S(v, G) : v \in V(G)\}$ is the *set of vertex stars of G*,

$\mathcal{NC}_k^*(G)$ is the set of non-separating cocircuit of matroid $M_k(G)$, and

$\mathcal{S}_k(G)$ is the set of vertex stars S of G such that $|S| \leq \rho_k^*(G)$, and so $\mathcal{S}_k(G)$ is the set of k-small vertex stars S of G.

We also remind that graphs $G = (V, E, \phi)$ and $G' = (V', E', \phi')$ with $E = E'$ are *strongly isomorphic* if there exists a bijection $\nu : V \to V'$ such that $\phi(e) = \{x, y\} \Leftrightarrow \phi'(e) = \{\nu(x), \nu(y)\}$.

Matroid M is called *k-circular* if there exists a graph G such that M is the k-circular matroid of G, i.e. $M = M_k(G)$. If $M_k(G) = M_k(G')$ implies that graphs G and G' are strongly isomorphic, then we say that G is *uniquely defined by* $M_k(G)$.

A natural problem $(WP)'_k$ is to describe all graphs G that are uniquely defined by $M_k(G)$. In this paper we establish some properties of graphs guaranteeing that the graphs are uniquely defined by their k-circular matroids.

It is easy to see that graphs G and G' with $E(G) = E(G')$ are strongly isomorphic if and only if $\mathcal{S}(G) = \mathcal{S}(G')$. For that reason the notion of a vertex star in a graph will play a central role in our discussion of the problem $(WP)'_k$.

We will distinguish between three types of vertices according to the size of their vertex stars in a graph G with respect to the rank $\rho_k^*(G)$ of matroid $M_k^*(G)$. We recall [6] that if

$M_k(G)$ is a connected matroid, then $\rho_k^*(G) = \Delta G - k + 1$.

Definition 2.1 Let x be a vertex in G, $k \geq 0$, $M_k(G)$ a connected matroid, and $\rho_k^*(G)$ the corank of $M_k(G)$. Then x is called *a k-small vertex of G* if $s(x, G) < \rho_k^*(G) + 1$, *a k-tight vertex of G* if $s(x, G) = \rho_k^*(G) + 1$, and *a k-big vertex of G* if $s(x, G) > \rho_k^*(G) + 1$.



Figure 1: Changes of the status of vertices when k increases from 0 to 5.



Figure 2: Changes of the status of vertices when k increases from 6 to 8.

3 Stars of G that are cocircuits in $M_k(G)$

We start with the following simple and useful observation.

Claim 3.1 *If K is a cocircuit of a matroid, then $|K| \leq \rho^*(M) + 1$.*

We remind that if $B \in \mathcal{B}_k(G)$ and $e \in B$, then $K(e, B)$ denotes the fundamental cocircuit of B rooted at e.

Claim 3.2 *Let G be a graph, $k \geq 1$, and $x \in V(G)$. Suppose that $M_k(G)$ is a connected matroid, B is a base of $M_k(G)$, $s(x, G\langle B\rangle) = 1$, and e is the edge incident to x in $G\langle B\rangle$. Then $S(x, G) = K(e, B)$, and so $S(x, G) \in \mathcal{C}_k^*(G)$.*

Proof (uses Theorems 4.6 and 4.7)
First, suppose that e is a loop in G. Since $s(x, G\langle B\rangle) = 1$, clearly, $G\langle e\rangle$ is a cycle component of $G\langle B\rangle$ with exactly one vertex x and one edge e. Then by Theorem 4.7, $K(e, B)\backslash e$ is the set of edges in $E\backslash B$ having x as an end vertex. Therefore $S(x, G) = K(e, B)$.

Now, suppose that e is not a loop in G. Then x is a leaf in $G\langle B\rangle$. By Theorem 4.6, again $S(x, G) = K(e, B)$. □

We remind the following known fact on matroids.

Claim 3.3 *Let $M = (E, \mathcal{I})$ be a matroid. Then*

($c1$) if $B \in \mathcal{B}(M)$ and $e \in E \setminus B = B^*$, then there exists a unique circuit $C = C(e, B)$ of M such that $e \in C \subseteq B \cup e$ (or, equivalently, such that $C \cap B^* = \{e\}$),

($c2$) similarly, if $B \in \mathcal{B}(M)$ and $e \in B$, then there exists a unique cocircuit $C^* = C^*(e, B)$ of M such that $e \in C^* \subseteq B^* \cup e$ (or, equivalently, such that $C^* \cap B = \{e\}$),

($c3$) $u \in C(e, B)\backslash e \Leftrightarrow (B\backslash u)\cup e \in \mathcal{B}(M)$ and similarly, $u \in C^*(e, B)\backslash e \Leftrightarrow (B\backslash e)\cup u \in \mathcal{B}(M)$, and

($c4$) for every $C \in \mathcal{C}(M)$ ($C^* \in \mathcal{C}(M)$) there exists $B \in \mathcal{B}$ and $e \in E \setminus B = B^*$ such that $C = C(e, B)$ (respectively, $e \in B$ such that $C^* = C^*(e, B)$).

Claim 3.4 *Let G be a graph, x a vertex of G, and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then the following are equivalent:*

($c1$) $S(x, G) \in \mathcal{C}_k^*(G)$ and

($c2$) for every edge e in $S(x, G)$ there exists $B \in \mathcal{B}_k(G)$ such that e is either a dangling edge at x or the edge of a loop component in $G\langle B\rangle$.

Proof (uses Theorems 4.6 and 4.7 and Claims 3.2 and 3.3)
First, we prove ($c1$) $\Rightarrow$ ($c2$). Since $S = S(x, G)$ is a cocircuit, by Claim 3.3 ($c3$), there exists $D \in \mathcal{B}_k(G)$ and $d \in D$ such that $S = K(d, D)$, and so $S \cap D = d$. Since S is a vertex star of G, d is either a dangling edge at x or the edge of a loop component in $G\langle D\rangle$. Let $e \in S = K(d, D)$. Then by Theorem 4.6 in case edge d is dangling and by Theorem 4.7 in case edge d is the edge of a loop component in $G\langle D\rangle$, we have: $B = (D \setminus d) \cup e \in \mathcal{B}_k(G)$. Clearly,

$K(e, B) = K(d, D) = S$. By the previous argument applied to B, e, and $S = K(e, B)$, e is either a dangling edge or the edge of a loop component in $G\langle B\rangle$.

Now, we prove $(c2) \Rightarrow (c1)$. Let $e \in S(x, G)$ and $B \in \mathcal{B}_k(G)$ such that e is either a dangling edge at x or the edge of a loop component in $G\langle B\rangle$. Then $s(x, G\langle B\rangle) = 1$. By Claim 3.2, $S(x, G) \in \mathcal{C}_k^*(G)$. □

We need the following three claims from [6].

Claim 3.5 (see Claim 4.4.11 in [6]) *Let $k \geq 1$. If $M_k(G)$ is a connected matroid, then $G \in \mathcal{G}_{\bowtie}$ and $\Delta G \geq k$.*

Theorem 3.6 (4.5.3 in [6]) GRAPH STRUCTURE OF A BASE OF $M_k(G)$ IN GRAPH G

Let G be a graph and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then the following are equivalent:

$(c1)$ $B \in \mathcal{B}_k(G)$ and

$(c2)$ $\Delta G\langle B\rangle = k - 1$, $V(G\langle B\rangle) = V(G)$ (i.e. B spans $V(G)$), and $\Delta A \geq 0$ for every component A of $G\langle B\rangle$ (i.e. $G\langle B\rangle$ has no tree component).

Corollary 3.7 (4.5.4 in [6]) *Let G be a graph and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then $\rho_k(G) = |V(G)| - 1 + k$ and $\rho_k^*(G) = |E(G)| - |V(G)| + 1 - k$.*

Claim 3.8 *Let G be a graph and $k \geq 1$. Suppose that*

$(a1)$ $M_k(G)$ is a connected matroid and

$(a2)$ x is a vertex of G such that $s(x, G) \leq \rho_k^*(G) + 1$, i.e. x is not a k-big vertex of G.

Then the following are equivalent:

$(c1)$ there exists $B \in \mathcal{B}_k(G)$ such that $x \notin V\lfloor G\langle B\rangle\rfloor$ and

$(c2)$ $Q \setminus x$ has a cycle, where Q is the component of G containing x.

Proof (uses Theorem 3.6, Corollary 3.7, and Claim 3.5)

(**p1**) We prove $(c1) \Rightarrow (c2)$. Suppose that $B \in \mathcal{B}_k(G)$ and $x \notin V\lfloor G\langle B\rangle\rfloor$. By Theorem 3.6, every component of $G\langle B\rangle$ has a cycle. Let A be the component of $G\langle B\rangle$ containing x. Since $x \notin V\lfloor G\langle B\rangle\rfloor$, clearly, A has a cycle C avoiding x. Obviously, Q contains A, and therefore also contains C.

(**p2**) We prove $(c2) \Rightarrow (c1)$. Since $M_k(G)$ is connected, by Claim 3.5, $G \in \mathcal{G}_{\bowtie}$. Since $G \in \mathcal{G}_{\bowtie}$, clearly, $G \setminus e$ has no tree component for every edge e in G. Then there exists a maximal subset Z of $S = S(x, G)$ such that $G' = G \setminus Z$ has no tree component and there is an edge p in $S \setminus Z$ incident to a component of $G \setminus x$ having a cycle. By assumption $(c2)$, such edge set Z exists and by the maximality of Z edge p is unique. Therefore $a \notin E\lfloor G'\rfloor$ for every $a \in S \setminus Z = S \cap E(G')$. Thus, $x \notin V\lfloor A'\rfloor$ for every $A' \in Cmp(G')$.

Since $G \setminus e$ has no tree component for every edge e in G, set Z is a proper subset of S. Then, clearly, $V(G') = V(G)$. Since $M_k(G)$ is a connected matroid, by Corollary 3.7, we have: $\rho_k^*(G) = |E(G)| - |V(G)| + 1 - k$. Since $s(x, G) \leq \rho_k^*(G) + 1$, we have:

$|E(G)|-s(x,G)+1-|V(G)| \geq k-1$. Since $|Z| < s(x,G)$, we have: $\Delta G' = |E(G')|-|V(G')| = |E(G)| - |Z| - |V(G)| \geq |E(G)| - s(x,G) + 1 - |V(G)| \geq k-1$.

Let $\mathcal{F}$ be the family of subgraphs F of G' such that $V(F) = V(G')$, $\Delta F \geq k-1$, and F has no tree component. Let $H \in \mathcal{F}$ be such that ΔH is minimum. Let $B = E(H)$. Since H is a subgraph of G', clearly $x \notin V\lfloor A \rfloor$ for every $A \in Cmp(H) = Cmp(G\langle B \rangle)$. We claim that $B \in \mathcal{B}_k(G)$. By Theorem 3.6, it is sufficient to prove that $\Delta H = k-1$.

First, suppose that every component of H has exactly one cycle. Then $\Delta H = 0$. Since $0 = \Delta H \geq k-1$ and k is a postive integer, clearly $k = 1$, and therefore $\Delta H = k-1$.

Now, suppose that H has a component D such that D has at least two cycles. Let e be an edge that belongs to a cycle in D. Let $H' = H \setminus e$. Then, clearly, $V(H') = V(G')$, H' has no tree component, and $\Delta H' = \Delta H - 1$. By the selection of H, graph $H' \notin \mathcal{F}$, and so $\Delta H' < k-1$. Therefore, $\Delta H = k-1$. □

Claim 3.9 (3.3.9 in [6]) *Let A be a connected graph with at least one cycle and $e \in E(A)$. Then*

($c0$) $e \in E\lfloor A \rfloor$ if and only if both end vertices of e belong to $\lfloor A \rfloor$,

($c1$) if $e \notin E\lfloor A \rfloor$, then $A \setminus e$ has two components and exactly one of them is a tree and the other component contains $\lfloor A \rfloor$,

($c2$) if A has one cycle and $e \in E\lfloor A \rfloor$, then $A \setminus e$ is a tree,

($c3$) if A has at least two cycles and $e \in E\lfloor A \rfloor$, then every component of $A \setminus e$ contains a cycle, and

($c4$) if $A \setminus e$ has two components and $v \in V\lfloor A \rfloor$, then the component of $A \setminus e$ containing v is not a tree.

Notation 3.10 Let G be a graph and $x \in V(G)$. If $G \setminus x$ has a cycle, then let G_x denote the union of all non-tree components of $G \setminus x$. If $G \setminus x$ has a tree component, then let G^x denote the union of all tree components of $G \setminus x$.

Claim 3.11 *Let G be a graph, $x \in V(G)$, and $k \geq 1$. Suppose that*

($a1$) $M_k(G)$ is a connected matroid and

($a2$) there exists $B \in \mathcal{B}_k(G)$ such that $x \notin V\lfloor G\langle B \rangle \rfloor$.

Then there exists $B' \in \mathcal{B}_k(G)$ such that

($c1$) T is a tree component of $G\langle B' \rangle \setminus x$ if and only if T is a tree component of $G \setminus x$,

($c2$) $G\langle B' \rangle \setminus V(G_x)$ is a tree, and

($c3$) there is exactly one edge in $G\langle B' \rangle$ from x to $V(G_x)$.

Proof (uses Theorem 3.6 and 4.6 and Claims 3.9 and 3.3)

Let $\mathcal{F} = \{B \in \mathcal{B}_k(G) : x \notin V\lfloor G\langle B \rangle \rfloor\}$. By assumption ($a2$) of our claim, $\mathcal{F} \neq \emptyset$. Let $B' \in \mathcal{F}$ be such that $s(x, G\langle B' \rangle)$ is minimum. Since $B' \in \mathcal{B}_k(G)$, by Theorem 3.6, B' spans $V(G)$ and $G\langle B' \rangle$ has no tree components. We claim that $G\langle B' \rangle$ satisfies conditions ($c1$)−($c3$) of our claim.

Since by $(a2)$, $x \notin V\lfloor G\langle B'\rangle\rfloor$, graph $G \setminus x$ has a cycle and G_x is defined.

Let A' be the component of $G\langle B'\rangle$ containing x, and so $x \notin V\lfloor A'\rfloor$. Since x is not incident to $\lfloor A'\rfloor$, clearly, $e \notin E\lfloor A'\rfloor$ for every $e \in S(x, B')$.

By Claim 3.9 $(c2)$, for every $e \in S(x, A')$ the graph $A' \setminus e$ consists of two components: one of these two components is a tree T_e and the other component U_e contains $\lfloor A'\rfloor$.

Let P be a minimal path in A' from x to $\lfloor A'\rfloor$. We claim that P is the only minimal path in A' from x to $\lfloor A'\rfloor$. Indeed, if there is another minimal path P' from x to $\lfloor A'\rfloor$, then the core of $P \cup P' \cup \lfloor A'\rfloor$ is a subgraph of A' containing $\lfloor A''\rfloor$ properly, contradicting the definition of $\lfloor A'\rfloor$. Obviously, $S(x, A')$ and $E(P)$ have exactly one edge in common, say f.

Suppose that there exists a tree component T of $G\langle B'\rangle \setminus x$. Then clearly, $T = T_e$ for some $e \in S(x, A') \setminus f$. We claim that T is a tree component of $G \setminus x$. Indeed, suppose not. Then there exists $g \in E \setminus B'$ such that g is incident to T and not incident to x. Therefore by Theorem 4.6, $g \in K(e, B')$. By Claim 3.3, $B'' = (B' \setminus e) \cup g$ is a k-base of G. Clearly, $x \notin V\lfloor G\langle B''\rangle\rfloor$, and so $B'' \in \mathcal{F}$. But $s(x, G\langle B''\rangle) < s(x, G\langle B'\rangle)$, contradicting the selection of B'.

Now, suppose that there exists a tree component T of $G \setminus x$. Then every cycle of G having vertices of T contains x. Since B' spans $V(G)$ and $G\langle B'\rangle$ has no tree components, there exists an edge $e \in B'$ incident to a vertex z of T and $V(G) \setminus V(T)$. Since T is a tree component of $G \setminus x$, edge e is incident to vertex x. Since T_e is a component of $G\langle B'\rangle \setminus e$, T_e is a subgraph of the component of $G \setminus x$ containing z. Therefore T_e is a subtree of T. By the argument in the previous paragraph T_e is a tree component of $G \setminus x$. Therefore $T_e = T$ and e is the only edge in B' having exactly one end vertex in T. Thus, our claims $(c1)$ and $(c2)$ are true.

Since T is a tree component of $G\langle B'\rangle \setminus x$ if and only if T is a tree component of $G \setminus x$, every vertex of $V(G_x)$ belongs to a non-tree component of $G\langle B'\rangle \setminus x$. If there are two edges in $G\langle B'\rangle$ from x to $V(G_x)$, then $x \in [G\langle B'\rangle]$, contradicting that $x \notin V\lfloor G\langle B'\rangle\rfloor$. Therefore f is the only edge in $G\langle B'\rangle$ from x to $V(G_x)$, and so $(c3)$ of our claim is true. □

Claim 3.12 *Let G be a graph and $k \geq 1$. Suppose that*

$(a1)$ *$M_k(G)$ is a connected matroid,*

$(a2)$ *there exists $B \in \mathcal{B}_k(G)$ and $x \in V(G)$ such that $x \notin V\lfloor G\langle B\rangle\rfloor$, and*

$(a3)$ *$G \setminus x$ has no tree component.*

Then there exists $B' \in \mathcal{B}_k(G)$ such that x is a leaf in $G\langle B'\rangle$.

Proof (uses Claims 3.11)

The assumptions $(a1)$ and $(a2)$ of our claim are the assumptions $(a1)$ and $(a2)$, respectively of Claim 3.11. Therefore there exists $B' \in \mathcal{B}_k(G)$ satisfying claims $(c1)-(c3)$ of Claim 3.11. Since $G \setminus x$ has no tree component, by Claim 3.11 $(c1)$, graph $G\langle B'\rangle \setminus x$ has no tree component. Therefore $V(G) = V(G_x) \cup \{x\}$. By Claim 3.11 $(c2)$, $G\langle B'\rangle \setminus V(G_x)$ is the isolated vertex x. By Claim 3.11 $(c3)$, there is exactly one edge in $G\langle B'\rangle$ from x to $V(G_x)$, and so x is a leaf in $G\langle B'\rangle$. □

From Claims 3.8 and 3.12, we have:

Claim 3.13 *Let G be a graph and $k \geq 1$. Suppose that*

(a1) $M_k(G)$ is a connected matorid,

(a2) x is a vertex of G such that $s(x, G) \leq \rho_k^*(G) + 1$, i.e. x is not a k-big vertex of G, and

(a3) $G \setminus x$ has no tree component.

Then exactly one of the following holds:

(c1) the component of G containing vertex x has no other vertex,

(c2) there exists $B \in \mathcal{B}_k(G)$ such that x is a leaf in $G\langle B\rangle$.

From Claims 3.4 and 3.13, we have:

Theorem 3.14 A CONDITION FOR A STAR TO BE A k-COCIRCUIT
Let G be a graph, $k \geq 1$, and $x \in V(G)$. Suppose that

(a1) $M_k(G)$ is a connected matroid and

(a2) $s(x, G) \leq \rho_k^*(G) + 1$, i.e. x is not a k-big vertex of G.

Then the following are equivalent:

(c1) $S(x, G) \in \mathcal{C}_k^*(G)$ and

(c2) $G \setminus x$ has no tree component.

Proof (uses Theorem 3.6, Corollary 3.7, and Claims 3.4, 3.8, and 3.13)

(p1) First, we prove (c1) $\Rightarrow$ (c2). Let $e \in S(x, G)$. Since $S(x, G) \in \mathcal{C}_k^*(G)$, by Claim 3.4, there exists $B \in \mathcal{B}_k(G)$ such that e is either a dangling edge or the edge of a loop component in $G\langle B\rangle$. By Theorem 3.6, every component in $G\langle B\rangle$ contains a cycle. Therefore every component of $G\langle B\rangle \setminus x$ has a cycle. Thus, $G \setminus x$ has no tree component.

(p2) Now, we prove (c2) $\Rightarrow$ (c1). If the component of G containing x has at least two vertices, by Claim 3.13, there exists $B \in \mathcal{B}_k(G)$ such that x is a leaf in $G\langle B\rangle$. Therefore, by Claim 3.4, $S = S(x, G) \in \mathcal{C}_k^*(G)$. So we assume that the component of G containing vertex x has no other vertex, and so every edge in $S(x, G)$ is a loop in G.

In what follows, the arguments we use are similar to those in (**p2**) of the proof of Claim 3.8.

Let $e \in S = S(x, G)$ and $Z = S \setminus e$. Let $G' = G \setminus Z$. Since $M_k(G)$ is a connected matroid, by Corollary 3.7, we have: $\rho_k^*(G) = |E(G)| - |V(G)| + 1 - k$. Since $s(x, G) \leq \rho_k^*(G) + 1$, we have: $|E(G)| - s(x, G) + 1 - |V(G)| \geq k - 1$. Since $|Z| < s(x, G)$, we have: $\Delta G' = |E(G')| - |V(G')| = |E(G)| - |Z| - |V(G)| \geq |E(G)| - s(x, G) + 1 - |V(G)| \geq k - 1$. Clearly $V(G') = V(G)$, $\Delta G' \geq k - 1$, and G' has no tree component.

Let $\mathcal{F}$ be the family of subgraphs F of G' such that $V(F) = V(G')$, $\Delta F \geq k - 1$, and F has no tree component. Let $H \in \mathcal{F}$ be such that ΔH is minimum. Let $B = E(H)$. We claim that $B \in \mathcal{B}_k(G)$. By Theorem 3.6, it is sufficient to prove that $\Delta H = k - 1$.

First, suppose that every component of H has exactly one cycle. Then $\Delta H = 0$. Since $0 = \Delta H \geq k - 1$ and k is a postive integer, clearly $k = 1$, and therefore $\Delta H = k - 1$.

Now, suppose that H has a component D such that D has at least two cycles. Let p be an edge that belongs to a cycle in D. Let $H' = H \setminus p$. Then clearly, $V(H') = V(G')$, H' has no tree component, and $\Delta H' = \Delta H - 1$. By the selection of H, graph $H' \notin \mathcal{F}$, and so $\Delta H' < k - 1$. Therefore, $\Delta H = k - 1$.

Thus, in both cases $\Delta H = k - 1$, and so $B \in \mathcal{B}_k(G)$. By definition of H and since the component of G containing vertex x has no other vertex, clearly e is the only edge of the component of $H = G\langle B\rangle$ containing x. By Claim 3.4, $S(x, G) \in \mathcal{C}_k^*(G)$. □

4 Stars of G and non-separating cocircuits of $M_k(G)$

We will use the following three facts from [6].

Theorem 4.1 (4.2.1 in [6]) A criterion for matroid $M_k(G)$ to be non-trivial

Let G be a graph, $F(G)$ the union of all tree components of G, and $k \geq 1$. Then the following are equivalent:
(a1) $\mathcal{C}_k(G) \neq \emptyset$,
(a2) $k \leq \Delta G + cmp(F(G))$, and
(a3) $M_k(G)$ is a non-trivial matroid.

Theorem 4.2 (4.4.5 in [6]) Graph description of connected matroid $M_k(G)$

Let $k \geq 2$. Then the following are equivalent:

(a1) $M_k(G)$ is a non-trivial matroid and $G \in \mathcal{G}_{\bowtie}$ and

(a2) $M_k(G)$ is a connected matroid.

Theorem 4.3 (4.4.6 in [6]) Graph description of connected matroid $M_1(G)$

The following are equivalent:

(a1) $M_1(G)$ is a non-trivial matroid and $G \in \mathcal{CG}_{\bowtie}$ and

(a2) $M_1(G)$ is a connected matroid.

Recall that a cocircuit K of a connected matroid M is *non-separating* if and only if matroid $M \setminus K$ is connected. Given a graph G, let $\mathcal{NC}_k^*(G)$ denote the set of non-separating cocircuit of $M_k(G)$.

Here is a graph description of non-separating cocircuits of $M_k(G)$ in terms of k and G.

Claim 4.4 *Let G be a graph, $K \subseteq E(G)$, and $k \geq 1$. Then the following are equivalent:*

(c1) $K \in \mathcal{NC}_k^*(G)$ and

(c2) $\Delta(G \setminus K) \geq k$ and

(c2.1) $G \setminus K \in \mathcal{CG}_{\bowtie}$ if $k = 1$ and

(c2.2) $G \setminus K \in \mathcal{G}_{\bowtie}$ if $k \geq 2$.

Proof (uses Theorem 4.1, 4.2, and 4.3)

By definition, $K \in \mathcal{NC}_k^*(G)$ if and only if $M_k(G) \setminus K$ is a connected matroid. Clearly, $M_k(G) \setminus K = M_k(G \setminus K)$. Therefore $K \in \mathcal{NC}_k^*(G)$ if and only if matroid $M_k(G \setminus K)$ is connected.

Suppose that $k = 1$. By Theorem 4.3 for $k = 1$, $M_1(G \setminus K)$ is connected if and only if $M_1(G)$ is non-trivial and $G \in \mathcal{CG}_{\bowtie}$.

Next, suppose that $k \geq 2$. By Theorem 4.2 for $k \geq 2$, $M_k(G \setminus K)$ is connected if and only if $M_k(G)$ is non-trivial and $G \in \mathcal{CG}_{\bowtie}$.

Now, if $G \setminus K \in \mathcal{G}_{\bowtie}$, then $G \setminus K$ has no tree components and by Theorem 4.1, $M_k(G)$ is non-trivial if and only if $\Delta(G \setminus K) \geq k$. Thus, $(c1) \Leftrightarrow (c2)$. □

We will distinguish between three possible types of (B, e)-cocircuits $K(e, B)$ depending on the structure of component A of $G\langle B\rangle$ containing edge e and on the position of edge e in A.

Definition 4.5 (4.7.1 in [6])

Let $B \in \mathcal{B}_k(G)$, $e \in B$, and $k \geq 1$. Then

$(t1)$ $K(e, B)$ is a (B, e)-*cocircuit in* $M_k(G)$ *of type 1*, if $e \notin E\lfloor A\rfloor$, where A is a component of $G\langle B\rangle$ containing edge e,

$(t2)$ $K(e, B)$ is a (B, e)-*cocircuit in* $M_k(G)$ *of type 2*, if $e \in E\lfloor A\rfloor$, where A is a unicyclic component of $G\langle B\rangle$ containing edge e, and

$(t3)$ $K(e, B)$ is a (B, e)-*cocircuit in* $M_k(G)$ *of type 3*, if $e \in E[A]$, where A is a component of $G\langle B\rangle$ that has at least two cycles and contains edge e.

Theorem 4.6 (4.7.2 in [6]) Graph description of rooted cocircuits of type 1

Let G be a graph and $k \geq 1$. Suppose that

$(a1)$ *$M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and*

$(a2)$ *$K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 1 (i.e. edge $e \notin E\lfloor A\rfloor$, where A is a component of $G\langle B\rangle$ containing edge e).*

Then exactly one of the two components of $A \setminus e$ is a tree T and $K(e, B) = K'(e, B) \cup e$, where $K'(e, B)$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(T)$.

Theorem 4.7 (4.7.3 in [6]) Graph description of rooted cocircuits of type 2

Let G be a graph and $k \geq 1$. Suppose that

$(a1)$ *$M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and*

$(a2)$ *$K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 2, (i.e. $e \in E\lfloor A\rfloor$, where A is a unicyclic component of $G\langle B\rangle$ containing edge e).*

Then $A \setminus e$ is a tree and $K(e, B) = K'(e, B) \cup e$, where $K'(e, B)$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(A \setminus e)$.

Theorem 4.8 (4.7.4 in [6]) GRAPH DESCRIPTION OF ROOTED COCIRCUITS OF TYPE 3

Let G be a graph and $k \geq 1$. Suppose that

(a1) $M_k(G)$ is a connected matroid, $B \in \mathcal{B}_k(G)$, and $e \in B$ and

(a2) $K(e, B)$ is a (B, e)-cocircuit in $M_k(G)$ of type 3, (i.e. $e \in E[A]$, where A is a component of $G\langle B\rangle$ that has at least two cycles and contains edge e).

Then $K(e, B) = (E \setminus B) \cup e$.

Claim 4.9 *Let G be a graph and $k \geq 1$. If $M_k(G)$ is a connected matroid, then $\mathcal{NC}_k^*(G) \subseteq S(G)$.*

Proof (uses Theorems 4.6, 4.7, and 4.8 and Claim 3.3)

Since $K \in \mathcal{C}_k^*(G)$, by Claim 3.3 $(c3)$, there exists $B \in \mathcal{B}_k(G)$ and $e \in B$ such that K is the (B, e)-cocircuit in $M_k(G)$.

First, suppose that K is the (B, e)-cocircuit in $M_k(G)$ of type 3. Then by Definition 4.5, $e \notin E\lfloor A\rfloor$, where A is a component of $G\langle B\rangle$ containing edge e. Hence by Theorem 4.8, $K = (E(G) \setminus B) \cup e$, and so $E(G) \setminus K \subset B$. Therefore $E(G) \setminus K \in \mathcal{I}_k(G)$. Thus, $M_k(G) \setminus K$ is not a connected matroid, contradicting $K \in \mathcal{NC}_k^*(G)$.

Next, suppose that K is the (B, e)-cocircuit in $M_k(G)$ of type 2. Then by Definition 4.5, $e \in E\lfloor A\rfloor$, where A is a unicyclic component of $G\langle B\rangle$ containing edge e. Hence by Theorem 4.7, $A \setminus e$ is a tree and $K \setminus e$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(A \setminus e)$. If A has at least two edges, then $A \setminus e$ is a tree component of $G \setminus K$ having at least one edge. Clearly, no edge of $A \setminus e$ belongs to a k-circuit of $G \setminus K$. Therefore $M_k(G \setminus K)$ is not a connected matroid, again contradicting $K \in \mathcal{NC}_k^*(G)$. Therefore e is the only edge of the component of $G\langle B\rangle$ and it is a loop incident to exactly one vertex, say x. Clearly, $S(x, G)$ is the set of edges in $E \setminus B$ incident to x, and so $K = S(x, G)$.

Finally, suppose that K is the (B, e)-cocircuit in $M_k(G)$ of type 1. Then by Definition 4.5, $e \in E[A]$, where A is a component of $G\langle B\rangle$ that has at least two cycles and contains edge e. Hence by Theorem 4.6, exactly one of the two components of $A \setminus e$ is a tree T and $K \setminus e$ is the set of edges in $E \setminus B$ having at least one end-vertex in $V(T)$. Therefore T is a tree component of $G \setminus K$. If $|V(T)| \geq 2$, then $|E(T)| \geq 1$ and no edge of T belongs to a k-circuit of $G \setminus K$. Therefore $M_k(G \setminus K)$ is not a connected matroid, contradicting $K \in \mathcal{NC}_k^*(G)$. Thus, T has exactly one vertex, say x. Clearly, is $S(x, G)$ is the set of edges in $E \setminus B$ incident to x, and so $K = S(x, G)$.

Thus, we have proved that if $K \in \mathcal{NC}_k^*(G)$, then there exists $x \in V(G)$ such that $K = S(x, G)$. □

Claim 4.10 *Let G be a graph, x a vertex of G, and $k \geq 1$. Suppose that*

(a1) $M_k(G)$ is a connected matroid and

(a2) $s(x, G) \leq \rho_k^*(G)$, i.e. x is a k-small vertex of G.

Then $M_k(G \setminus x)$ is a non-trivial matroid.

Proof (uses Theorem 4.1 and Corollary 3.7)

First, we prove that $M_k(G \setminus x)$ is a non-trivial matroid. Since $M_k(G)$ is a connected matroid, by Corollary 3.7, we have: $\rho_k^*(G) = |E(G)| - |V(G)| + 1 - k$. Since $s(x, G) \leq \rho_k^*(G)$, we have: $k \leq (|E(G)| - s(x, G)) - (|V(G)| - 1) = |E(G \setminus x)| - |V(G \setminus x)| = \Delta(G \setminus x)$. Therefore $\Delta(G \setminus x) \geq k$. Therefore by Theorem 4.1, $M_k(G \setminus x)$ is a non-trivial matroid. □

Claim 4.11 *Let G be a graph, x a vertex of G, and $k \geq 1$. Suppose that*

(a1) $M_k(G)$ is a connected matroid,

(a2) $s(x, G) \leq \rho_k^*(G)$, i.e. x is a k-small vertex of G, and

(a3) $G \setminus x \in \mathcal{CG}_{\bowtie}$ if $k = 1$ and $G \setminus x \in \mathcal{G}_{\bowtie}$ if $k \geq 2$.

Then $S(x, G) \in \mathcal{NC}_k^*(G)$.

Proof (uses Theorems 3.14, 4.2, and 4.3 and Claim 4.10)

By assumption $(a3)$, $G \setminus x$ has no tree components. Therefore by Theorem 3.14, $S = S(x, G) \in \mathcal{C}_k(G)$. Clearly, $M_k(G) \setminus S = M_k(G \setminus x)$. We need to prove that $M_k(G \setminus x)$ is a connected matroid. By Claim 4.10, $M_k(G \setminus x)$ is a non-trivial matroid.

Suppose that $k = 1$. Then since $G \setminus x \in \mathcal{CG}_{\bowtie}$, by Theorem 4.3, $M_k(G \setminus x)$ is a connected matroid, and so $S(x, G) \in \mathcal{NC}_k^*(G)$.

Now, suppose that $k \geq 2$. Then since $G \setminus x \in \mathcal{G}_{\bowtie}$, by Theorem 4.2, $M_k(G \setminus x)$ is a connected matroid, and so $S(x, G) \in \mathcal{NC}_k^*(G)$. □

Theorem 4.12 Graph structure of non-separating k-cocircuits

Let G be a graph and $k \geq 1$. Suppose that $M_k(G)$ is a connected matroid. Then the following are equivalent:

(c1) $K \in \mathcal{NC}_k^*(G)$ and

(c2) there exists $x \in V(G)$ such that $K = S(x, G)$, $s(x, G) \leq \rho_k^*(G)$, and

(c2.1) $G \setminus x \in \mathcal{CG}_{\bowtie}$ if $k = 1$ and

(c2.2) $G \setminus x \in \mathcal{G}_{\bowtie}$ if $k \geq 2$.

Proof (uses Theorem 4.2 and 4.3 and Claims 4.9 and 4.11)

By Claim 4.11, $(c2) \Rightarrow (c1)$. We prove $(c1) \Rightarrow (c2)$. Since $K \in \mathcal{NC}_k^*(G)$, by Claim 4.9, there exists $x \in V(G)$ such that $K = S(x, G)$. Since $K = S(x, G) \in \mathcal{C}_k^*(G)$, clearly $s(x, G) \leq \rho_k^*(G) + 1$. However, if $s(x, G) = \rho_k^*(G) + 1$, then K contains a cobase of $M_k(G)$, and so $E(G) \setminus K$ is a subset of a base of $M_k(G)$. Then $M_k(G \setminus x)$ has no k-circuit and therefore is not connected, contradicting $K \in \mathcal{NC}_k^*(G)$. Thus, $s(x, G) \leq \rho_k^*(G)$.

Since $K = S(x, G)$, we have: $M_k(G \setminus K) = M_k(G \setminus x)$. Since $K \in \mathcal{NC}_k^*(G)$, matroid $M_k(G \setminus x)$ is connected.

Suppose that $k = 1$. Since $M_1(G \setminus x)$ is connected, by Theorem 4.3, $G \setminus x \in \mathcal{CG}_{\bowtie}$.

Now, suppose that $k \geq 2$. Since $M_k(G \setminus x)$ is connected, by Theorem 4.2, $G \setminus x \in \mathcal{G}_{\bowtie}$. □

Let $\mathcal{S}_k(G)$ denote the set of vertex stars S of G such that $|S| \leq \rho_k^*(G)$.

It is known [9] that $\mathcal{NC}^*(M(G)) = \mathcal{S}(G)$, where $M(G)$ is the cycle matroid of graph G. The next theorem is an analog of the above fact for k-circular matroids $M_k(G)$.

Theorem 4.13 Let G be a 3-connected graph, $k \geq 1$, and $|E(G)| - |V(G)| \geq k$. Then $\mathcal{NC}^*(M_k(G)) = \mathcal{S}_k(G)$.

Proof (uses Theorems 4.1, 4.2, 4.3, and 4.12)

Since $|E(G)| - |V(G)| \geq k$, by Theorem 4.1, $M_k(G)$ is a non-trivial matroid. Since G is 3-connected, clearly $G \in \mathcal{G}_{\bowtie}$. Then by Theorem 4.2 for $k \geq 2$ and Theorem 4.3 for $k = 1$, matroid $M_k(G)$ is connected. Therefore by Theorem 4.12, $\mathcal{NC}^*(M_k(G)) \subseteq \mathcal{S}_k(G)$.

Since G is 3-connected, clearly $G \setminus x \in \mathcal{G}_{\bowtie}$ and $G \setminus x$ is a connected graph for every $x \in V(G)$. Therefore again by Theorem 4.12, $\mathcal{S}_k(G) \subseteq \mathcal{NC}^*(M_k(G))$.

Thus, $\mathcal{NC}^*(M_k(G)) = \mathcal{S}_k(G)$. □

5 Uniquely representable matroids with $\mathcal{S}(G) \subseteq \mathcal{NC}^*(G)$

Recall that a graph G is uniquely defined by $M_k(G)$ if $M_k(G) = M_k(G')$ implies that graphs G and G' are strongly isomorphic.

Claim 5.1 [9] *Let G be a graph with $|V(G)| \geq 4$. Suppose that $M(G)$ is a connected matroid. Then the following are equivalent:*

$(c1)$ *G is uniquely defined by $M(G)$ and*

$(c2)$ *$\mathcal{S}(G) \subseteq \mathcal{NC}^*(G)$.*

In this part we will describe some results on matroids $M_k(G)$ analogous to the implication $(c2) \Rightarrow (c1)$ in Claim 5.1 about matroid $M(G)$.

We start with the following useful fact.

Claim 5.2 *Let G and G' be graphs and $k \geq 1$. Suppose that*

$(a1)$ *$M_k(G)$ is a connected matroid,*

$(a2)$ *x is a vertex of G such that $S(x, G) \in \mathcal{NC}^*_k(G)$, and*

$(a3)$ *$M_k(G) = M_k(G')$.*

Then $S(x, G) \in \mathcal{S}(G')$.

Proof (uses Theorem 4.12 and Claim 4.9)

Since $M_k(G) = M_k(G')$, clearly $\mathcal{NC}^*_k(G) = \mathcal{NC}^*_k(G')$. Since $S(x, G) \in \mathcal{NC}^*_k(G)$, we have: $S(x, G) \in \mathcal{NC}^*_k(G')$. Since $M_k(G')$ is a connected matroid, by Claim 4.9, there exists $x' \in V(G')$ such that $S(x, G) = S(x', G')$. Thus, $S(x, G) \in \mathcal{S}(G')$. □

Claim 5.3 *Let G and G' be graphs and $k \geq 1$. Suppose that*

$(a1)$ *$M_k(G)$ is a connected matroid and*

$(a2)$ *$M_k(G) = M_k(G') := M_k$.*

Then $|V(G)| = |V(G')|$.

Proof (uses Corollary 3.7)

Since $M_k(G) = M_k(G')$, we have: $\rho_k(G) = \rho_k(G')$. Since M_k is a connected matroid, by Corollary 3.7, $\rho_k(G) = |V(G)| - 1 + k$ and $\rho_k(G') = |V(G')| - 1 + k$. Therefore $|V(G)| = |V(G')|$. □

Now, we are ready to prove our first results on graphs uniquely defined by their k-circular matroids analogous to implication $(c2) \Rightarrow (c1)$ in Claim 5.1 about cycle matroid $M(G)$ of graph G.

We need the following known fact.

Claim 5.4 [9] *Let G and G' be graphs and $E(G) = E(G')$. Then G and G' are strongly isomorphic if and only if $\mathcal{S}(G) = \mathcal{S}(G')$.*

Theorem 5.5 A condition for $M_k(G)$ to uniquely define graph G

Let G and G' be graphs and $k \geq 1$. Suppose that

$(a1)$ *$M_k(G)$ is a connected matroid,*

$(a2)$ *$\mathcal{S}(G) \subseteq \mathcal{NC}_k^*(G)$, and*

$(a3)$ *$M_k(G) = M_k(G')$.*

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Claims 5.4, 5.2, and 5.3)

Since $S(x, G) \in \mathcal{NC}_k^*(G)$ for every $x \in V(G)$, by Claim 5.2, $\mathcal{S}(G) \subseteq \mathcal{S}(G')$. Now, by Claim 5.3, $|V(G)| = |V(G')|$, and therefore $|\mathcal{S}(G)| = |\mathcal{S}(G')|$. It follows that $\mathcal{S}(G) = \mathcal{S}(G')$ and by Claim 5.4, G and G' are strongly isomorphic. □

It is known [15] (see also [9]) that multi-3-connected graphs G are uniquely defined by $M(G)$, where $M(G)$ is the cycle matroid of graph G. The next theorem is an analog of the above fact for k-circular matroids $M_k(G)$.

Theorem 5.6 A condition for a graph G to be uniquely defined by $M_k(G)$

Let G and G' be graphs and $k \geq 2$. Suppose that

$(a1)$ *$\Delta(G) \geq k$ and $G \in \mathcal{G}_{\bowtie}$,*

$(a2)$ *$\Delta(G \setminus x) \geq k$ and $G \setminus x \in \mathcal{G}_{\bowtie}$ for every vertex x in G, and*

$(a3)$ *$M_k(G') = M_k(G)$.*

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Theorems 4.1, 4.2, 4.12, and 5.5)

By our assumption $(a1)$ and Theorems 4.1 and 4.2, we have: $M_k(G)$ is a connected matroid. Since $M_k(G)$ is a connected matroid, by Corollary 3.7, we have:

$\rho_k^*(G) = |E(G)| - |V(G)| + 1 - k$.

Since $\Delta(G \setminus x) \geq k$ for every vertex x in G, we have:

$\Delta(G \setminus x) = |E(G \setminus x)| - |V(G \setminus x)| = (|E(G)| - s(x, G)) - (|V(G)| - 1) \geq k$

for every vertex x in G. Therefore $\rho_k^*(G) \geq s(x,G)$. By our assumption $(a2)$, $G \setminus x \in \mathcal{G}_{\bowtie}$ for every vertex x in G. Since $s(x,G) \leq \rho_k^*(G)$ for every vertex x in G, we have by Theorem 4.12: $S(x,G) \in \mathcal{NC}_k^*(G)$ for every $x \in V(G)$. Now, by Theorem 5.6, $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic. □

Theorem 5.7 A condition for a graph G to be uniquely defined by $M_1(G)$

Let G and G' be graphs. Suppose that

$(a1)$ $\Delta(G) \geq 1$ *and* $G \in \mathcal{CG}_{\bowtie}$,

$(a2)$ $\Delta(G \setminus x) \geq 1$ *and* $G \setminus x \in \mathcal{CG}_{\bowtie}$ *for every vertex x in G, and*

$(a3)$ $M_1(G') = M_1(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Theorems 4.1, 4.3, 4.12, and 5.6)

By our assumption $(a1)$ and Theorems 4.1 and 4.3, we have: $M_1(G)$ is a connected matroid. Since $M_1(G)$ is a connected matroid, by Corollary 3.7, we have: $\rho_k^*(G) = |E(G)| - |V(G)|$. Since $\Delta(G \setminus x) \geq 1$ for every vertex x in G, we have:

$$\Delta(G \setminus x) = |E(G \setminus x)| - |V(G \setminus x)| = (|E(G)| - s(x,G)) - (|V(G)| - 1) \geq 1$$

for every vertex x in G. Therefore $\rho_k^*(G) \geq s(x,G)$. By our assumption $(a2)$, $G \setminus x \in \mathcal{CG}_{\bowtie}$ for every vertex x in G. Since $s(x,G) \leq \rho_1^*(G)$ for every vertex x in G, we have by Theorem 4.12: $S(x,G) \in \mathcal{NC}_1^*(G)$ for every $x \in V(G)$. Now, by Theorem 5.6, $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic. □

Theorem 5.8 *Let G and G' be graphs and $k \geq 1$.*

Suppose that

$(a1)$ *G is a 3-connected graph,*

$(a2)$ $\mathcal{C}_k(G) \neq \emptyset$,

$(a3)$ $s(x,G) \leq \rho_k^*(G)$ *for every vertex x of G, and*

$(a4)$ $M_k(G') = M_k(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Theorems 4.1, 4.2, 4.3, 5.6, and 5.7)

Since G is a 3-connected graph, $G \in \mathcal{CG}_{\bowtie}$. Since $\mathcal{C}_k(G) \neq \emptyset$, by Theorem 4.1, matroid $M_k(G)$ is non-trivial. Therefore, by Theorems 4.2 for $k \geq 2$ and 4.3 for $k = 1$, $M_k(G)$ is a connected matroid. Clearly, all assumptions of Theorems 5.6 and 5.7 hold. Thus, $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic. □

Remark 5.9 Theorems 5.6 - 5.8 describe various conditions that guarantee a unique graph representation of some k-circular matroids.

6 Uniquely representable matroids with $\mathcal{S}(G) \not\subseteq \mathcal{NC}^*(G)$

Now, we will describe some results on matroids $M_k(G)$ illustrating a new phenomenon when implication $(c1) \Rightarrow (c2)$ in Claim 5.1 does not hold if matroid $M(G)$ is replaced by $M_k(G)$.

Theorem 6.1 *Let G and G' be graphs and $k \geq 1$.*

Suppose that

(a1) both graphs G and G' have no loops,

(a2) $M_k(G)$ is a connected matroid,

(a3) G has exactly one vertex v such that $S(v, G) \notin \mathcal{NC}_k^*(G)$, and

(a4) $M_k(G') = M_k(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Claims 5.4, 5.2, and 5.3)

If x is a vertex of G distinct from v, then by Claim 5.2, $S(x, G) \in \mathcal{S}(G')$. Then $\mathcal{S}(G) \setminus \{S(v, G)\} \subseteq \mathcal{S}(G')$. By Claim 5.3, $|V(G)| = |V(G')|$, and therefore $|\mathcal{S}(G)| = |\mathcal{S}(G')|$. Therefore only one vertex, say v', in G' is such that $S(v', G') \notin \mathcal{S}(G) \setminus \{S(v, G)\}$. Then $\mathcal{S}(G') \setminus \{S(v', G')\} = \mathcal{S}(G) \setminus \{S(v, G)\}$. Thus, it remains to prove that $S(v, G) = S(v', G')$.

Let $e \in S(v', G')$. Then clearly, e belongs to at most one element of $\mathcal{S}(G') \setminus \{S(v', G')\}$. Since $\mathcal{S}(G') \setminus \{S(v', G')\} = \mathcal{S}(G) \setminus \{S(v, G)\}$, we have: e belongs to at most one element of $\mathcal{S}(G) \setminus \{S(b, G)\}$. Since G has no loops, e is incident to b. Hence $S(v', G') \subseteq S(v, G)$.

Since both G and G' have no loops, by the arguments similar to those above, $S(v, G) \subseteq S(v', G')$. Thus, $S(v, G) = S(v', G')$. Hence $\mathcal{S}(G) = \mathcal{S}(G')$, and by Claim 5.4, G and G' are strongly isomorphic. □

Theorem 6.2 Another condition for $M_k(G)$ to uniquely define graph G

Let G and G' be graphs and $k \geq 1$. Suppose that

(a1) $M_k(G)$ is a connected matroid,

(a2) $S(x, G) \in \mathcal{NC}_k^*(G)$ for every vertex x of G except for one vertex v,

(a3) if p is a loop in G, then p is incident to v,

(a4) v is not a k-big vertex of G and $G \setminus v$ has no tree component, and

(a5) $M_k(G') = M_k(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Proof (uses Theorems 3.6, 3.14 and 6.1 and Claims 3.4, 5.2, and 5.3)

Our first step in proving our theorem is to prove that there exists a bijection $\alpha : V(G) \to V(G')$ such that $S(x, G) = S(\alpha(x), G')$ for every $x \in V(G) \setminus v$.

If x is a vertex of G distinct from v, then by Claim 5.2, $S(x, G) \in \mathcal{S}(G')$, and so $\mathcal{S}(G) \setminus \{S(v, G)\} \subseteq \mathcal{S}(G')$. Therefore for $x \in V(G) \setminus v$ there exists a unique vertex x' such that $S(x', G') = S(x, G)$. In this case we put $\alpha(x) = x'$. Since $M_k(G)$ is a connected matroid and $M_k(G') = M_k(G)$, by Claim 5.3, $|V(G)| = |V(G')|$, and therefore $|\mathcal{S}(G)| = |\mathcal{S}(G')|$. Hence only one vertex, say v', in G' is such that $S(v', G') \notin \mathcal{S}(G) \setminus \{S(v, G)\}$. Clearly, $v' \neq \alpha(x) = x'$ for every $x \in V(G) \setminus v$. We put $v' = \alpha(v)$. Clearly, α is a bijection from $V(G)$ to $V(G')$ and $S(x', G') = S(x, G)$ for every $x \in V(G) \setminus v$.

Now, we will prove that α is a strong isomorphism from G to G'. Obviously, it is sufficient to prove that $S(v, G) = S(\alpha(v), G') = S(v', G')$.

(**p1**) First, suppose that G has no loops. By Theorem 6.1, it is sufficient to prove that G'

has no loops. Suppose not. Then G' has a loop, say e. Since G has no loops, e is not a loop in G, and so e is incident to two distinct vertices in G, say x_1 and x_2. We claim that $v \in \{x_1, x_2\}$. Indeed, suppose not. Then $v \notin \{x_1, x_2\}$. Therefore $e \in S(x_1, G) \cap S(x_2, G)$, and so $e \in S(x'_1, G') \cap S(x'_2, G')$. Since $x'_1 = \alpha(x_1) \neq \alpha(x_2) = x'_2$, e is not a loop in G', a contradiction. Thus, e in G is incident to v and to another vertex of G, say z.

Since $e \in S(z, G) = S(z', G')$ and e is a loop in G', clearly, edge e is a loop at $z' = \alpha(z)$ in G'. Now, by assumption $(a4)$ of our theorem, $|S(v, G)| \leq \rho^*_k(G) + 1$ and $G \setminus v$ has no tree component. By Theorem 3.14, $S(v, G) \in \mathcal{C}^*_k(G)$. Therefore, by Claim 3.4, there exists $B \in \mathcal{B}_k(G)$ such that e is a dangling edge at v in $G\langle B\rangle$. Since $M_k(G') = M_k(G)$, clearly, $B \in \mathcal{B}_k(G)$. Since G has no loops and $S(x', G') = S(x, G)$ for every $x \in V(G) \setminus v$, we have: $G\langle B\rangle \setminus v = G'\langle B\rangle \setminus v'$. Therefore v' is an isolated vertex in $G'\langle B\rangle$, and so B does not span $V(G')$, contradicting Theorem 3.6.

(**p2**) Now, suppose that G has a loop. Since $S(x', G') = S(x, G)$ for every $x \in V(G) \setminus v$, an edge p is a loop at v in G if and only if p is a loop at v' in G'.

First, suppose that $e \in S(v, G)$ and e is not a loop in G. By our assumption $(a4)$ and Theorem 3.14, $S(v, G) \in \mathcal{C}^*_k(G)$. Therefore, by Theorem 3.4, there exists $B \in \mathcal{B}_k(G)$ such that e is a dangling edge at v in $G\langle B\rangle$. Clearly, from our assumption $(a3)$ and equality $S(x', G') = S(x, G)$ for every $x \in V(G) \setminus v$ we have: $G\langle B\rangle \setminus v = G'\langle B\rangle \setminus v'$. If e is not incident to v' in G', then B does not span $V(G')$, contradicting Theorem 3.6. Thus, $e \in S(v', G')$.

Finally, suppose that $e \in S(v', G')$ and e is not a loop in G'. Since $S(x', G') = S(x, G)$ for every $x \in V(G) \setminus v$, clearly, e belongs to at most one element of $\mathcal{S}(G) \setminus \{S(v, G)\}$. By our assumption $(a3)$, edge e is incident to v in G. Therefore $e \in S(v, G)$. □

Using Theorem 6.2 and the arguments similar to those in the proof of Theorem 5.6, one can prove the following theorem.

Theorem 6.3 Another condition for G to be uniquely defined by $M_k(G)$

Let G and G' be graphs and $k \geq 2$. Suppose that

$(a1)$ *$\Delta(G) \geq k$ and $G \in \mathcal{G}_{\bowtie}$,*

$(a2)$ *$\Delta(G \setminus x) \geq k$ and $G \setminus x \in \mathcal{G}_{\bowtie}$ for every vertex x in G except for one vertex v,*

$(a3)$ *if p is a loop in G, then p is incident to v,*

$(a4)$ *v is not a k-big vertex of G and $G \setminus v$ has no tree component, and*

$(a5)$ *$M_k(G') = M_k(G)$.*

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Using Theorem 6.2 and the arguments similar to those in the proof of Theorem 5.7, one can prove the following theorem.

Theorem 6.4 Another condition for G to be uniquely defined by $M_1(G)$

Let G and G' be graphs. Suppose that

$(a1)$ *$\Delta(G) \geq 1$ and $G \in \mathcal{CG}_{\bowtie}$,*

$(a2)$ *$\Delta(G \setminus x) \geq 1$ and $G \setminus x \in \mathcal{CG}_{\bowtie}$ for every vertex x in G except for one vertex v,*

$(a3)$ *if p is a loop in G, then p is incident to v,*

(*a*4) *v is not a 1-big vertex of G and $G \setminus v$ has no tree component, and*
(*a*5) $M_1(G') = M_1(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Using Theorems 4.1, 4.2, 4.3, 6.3, and 6.4 and the arguments similar to those in the proof of Theorem 5.8, one can prove the following theorem.

Theorem 6.5 *Let G and G' be graphs, $t \in V(G)$, and $k \geq 1$. Suppose that*
(*a*1) *graph G is 3-connected,*
(*a*2) $\mathcal{C}_k(G) \neq \emptyset$,
(*a*3) $s(t, G) = \rho_k^*(G) + 1$ *and* $s(x, G) \leq \rho_k^*(G)$ *for every vertex x of G distinct from t, and*
(*a*4) $M_k(G') = M_k(G)$.

Then $\mathcal{S}(G) = \mathcal{S}(G')$, i.e. G and G' are strongly isomorphic.

Claim 6.6 (2.3.2 in [6])

Let G be a graph with $v(G) = v$ and $e(G) = e$. Suppose that G is 3-connected and has a vertex x such that $d(x, G) > e - v$. Then G is the wheel with center x, and so $d(x, G) = v - 1$ and $d(z, G) = 3$ for every $z \in V(G) \setminus x$.

From Claim 6.6 and Theorems 5.8 and 6.5 we have, in particular:

Theorem 6.7 *Every 3-connected graph G not isomorphic to a complete graph on 4 vertices is uniquely defined by its bicircular matroid $M_1(G)$.*

Another proof of Theorem 6.7 can be obtained using some results in [2].



$$K \not\approx^s K' \Leftrightarrow M_0(K) \neq M_0(K')$$

$$K \not\approx^s K' \text{ but } M_1(K) = M_1(K')$$

$$K \not\approx L \text{ but } M_1(K) = M_1(L)$$

Figure 3: Graph K is not uniquely defined by matroid $M_1(K)$.

Theorem 6.8 When is a 3-connected graph G uniquely defined by $M_k(G)$?

Let G be a graph, $k \geq 1$, and $M_k(G)$ the k-circular matroid of G. Suppose that G is 3-connected. Then the following are equivalent:

($c1$) *G is uniquely defined by $M_k(G)$ and*

($c2$) *$|E(G)| - |V(G)| \geq k$ and either every vertex of G is k-small or every vertex of G is k-small except for one which is k-tight.*

The implication ($c2$) $\Rightarrow$ ($c1$) of Theorem 6.8 follows immediately from Theorems 5.8 and 6.5. In our next paper we will describe some graph operations that will provide non-isomorphic graphs with the same k-circular matroid. Implication ($c1$) $\Rightarrow$ ($c2$) in Theorem 6.8 will follow from those results.

Theorem 6.8 is a natural extension of the classical Whitney's matroid-isomorphism theorem on the cycle matroid of a 3-connected graph [15] (see also [9]).

References

[1] J. A. Bondy and U. S. R. Murty, *Graph Theory*, 3rd Corrected Printing, GTM 244, Springer-Verlag, New York, 2008.

[2] C. R. Coulard, J. G. del Greco, D. K. Wagner, Representations of bicircular matroids, *Discrete Applied Mathematics* **32** (1991) 223-240.

[3] R. Diestel, *Graph Theory*, Springer-Verlag, New York, 2000.

[4] R. Halin and H. A. Jung, Note on isomorphisms of graphs, *J. London Math.* Sec. **42** (1967) 254-256.

[5] R. L. Hemminger, H. A. Jung, A. K. Kelmans, On 3–skein isomorphisms of a graph, *Combinatorica* **2** (4), (1982) 373-376.

[6] J. F. De Jesús and A. K. Kelmans, K-circular matroids of graphs, arXiv:1508.05364 (2015).

[7] A. K. Kelmans, On semi-isomorphisms and semi-dualities of graphs, *Graphs and Combinatorics* **10** (1994) 337–352.

[8] A. K. Kelmans, On 3-skeins in a 3-connected graph. *Studia Scientarum Mathematicarum Hungarica* **22** (1987) 265-273.

[9] A. K. Kelmans, The concept of a vertex in a matroid, the non-separating cycles and a new criterion for graph planarity. In *Algebraic Methods in Graph Theory*, Vol. 1, Colloq. Math. Soc. János Bolyai, (Szeged, Hungary, 1978) North–Holland **25** (1981) 345-388.

[10] J. G. Oxley, *Matroid Theory*, Oxford University Press, 2006.

[11] D. K. Wagner, Connectivity in bicircular matroids, *J. Combinatorial Theory* **B 39** (1985) 308-324.

[12] D. J. A. Welsh, *Matroid Theory*, Academic Press, London, 1976.

[13] H. Whitney, On the abstract properties of linear dependence, *Amer. J. Math.* **57** (1935) 509-533.

[14] H. Whitney, 2-isomorphic graphs, *Amer. J. Math.* **55** (1933) 245-254.

[15] H. Whitney, Congruent graphs and the connectivity of graphs, *Amer. J. Math.* **54** (1932) 159-168.

[16] H. Whitney, Non-separable and planar graphs, *Amer. J. Math.* **34** (1932) 339-362.